

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Sheep anti Human alpha-1 antichymotrypsin, Clone: [Polyclonal], Monoclonal NMB-SHAHU/ACHY

| | Sheep anti Human alpha-1 antichymotrypsin, Clone: [Polyclonal], |
|----------------------------|---|
| Article Name | Monoclonal |
| Biozol Catalog Number | NMB-SHAHU/ACHY |
| Supplier Catalog Number | ShAHu/Achy |
| Alternative Catalog Number | NMB-SHAHU/ACHY |
| Manufacturer | NordicMubio |
| Host | Sheep |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to alpha-1 antichymotrypsin. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against human plasma a single precipitin line is obtained and shows a reaction of f |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| NCBI | 12 |
| UniProt | P01011 |
| Buffer | Delipidated, heat inactivated, lyophilized, stable antiserum. No preservative added. Total protein and IgG concentration are comparable to those of pooled normal sheep serum. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. |

| Source | Alpha-1 antichymotrypsin is a protease inhibitor with a molecular weight of 68.000. Its concentration in fetal human serum is low at birth, but the concentration increases rapidly to normal plasma levels of about is 4.5 μ g/ml. Alpha-1 antichymotrypsin is |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized, stable antiserum. No preservative added. Total protein and IgG concentration are comparable to those of pooled normal sheep serum. |
| Application Notes | Precipitation assays. In immunoelectrophoresis in agarose-plates use 2 μ l human plasma or equivalent against 120 μ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 μ l antiserum in 3 mm diameter center well and 2 μ l plasma sa |