

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Swine anti Goat IgA (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAG/IGA(FC)

Article Name	Swine anti Goat IgA (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal
Biozol Catalog Number	NMB-SWAG/IGA(FC)
Supplier Catalog Number	SwAG/IgA(Fc)
Alternative Catalog Number	NMB-SWAG/IGA(FC)
Manufacturer	NordicMubio
Host	Porcine
Category	Antikörper
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgA molecule as tested in immunoelectrophoresis and radial immunodiffusion. In immunoelectrophoresis using various antiserum concentrations against normal goat serum only IgA is prec
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added. Reconstitute the lyophilized
Source	Highly purified secretory IgA isolated from pooled goat milk. Freunds complete adjuvant is used in the first step of the immunization procedure.

Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added.
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l serum or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 $\mu$ l antiserum in 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and serially d