

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Swine anti Horse IgG (heavy and light chains), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAHO/IGG(H+L)

Article Name	Swine anti Horse IgG (heavy and light chains), Clone: [Polyclonal], Porcine, Monoclonal
Biozol Catalog Number	NMB-SWAHO/IGG(H+L)
Supplier Catalog Number	SwAHo/IgG(H+L)
Alternative Catalog Number	NMB-SWAHO/IGG(H+L)
Manufacturer	NordicMubio
Host	Porcine
Category	Antikörper
Product Description	The reactivity of the antiserum is directed to the Fc and Fab subunits of both subclasses IgG1 and IgG2. Reactivity with other subclasses of horse IgG has not been tested. The antiserum reacts with both Fc and Fab portions of polyclonal IgG. It inclu
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled swine se
Source	Purified normal IgG isolated from pooled horse serum. Freunds complete adjuvant is used in the first step of the immunization procedure.

Formula	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled swine se
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samples (neat and