

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Swine anti Horse Ig light chains kappa and lambda (free and bound), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAHO/L(SD+HD)

| Article Name | Swine anti Horse Ig light chains kappa and lambda (free and bound), Clone: [Polyclonal], Porcine, Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NMB-SWAHO/L(SD+HD) |
| Supplier Catalog Number | SwAHo/L(SD+HD) |
| Alternative Catalog Number | NMB-SWAHO/L(SD+HD) |
| Manufacturer | NordicMubio |
| Host | Porcine |
| Category | Antikörper |
| Product Description | As tested in immunoelectrophoresis and double radial immunodiffusion against pooled normal horse serum and purified IgG fractions and light chain preparations, the antiserum reacts with determinants exposed on the surface (SD) and the Fab portion of |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled swine se |
| Source | Purified equine light chains isolated from polyclonal horse IgG. Freunds complete adjuvant is used in the first step of the immunization procedure. |

| Formula | Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled swine se |
|-------------------|---|
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samples (neat and |