

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Swine anti Human IgG (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAHU/IGG(FC)

Article Name	Swine anti Human IgG (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal	
Biozol Catalog Number	NMB-SWAHU/IGG(FC)	
Supplier Catalog Number	SwAHu/lgG(Fc)	
Alternative Catalog Number	NMB-SWAHU/IGG(FC)	
Manufacturer	NordicMubio	
Host	Porcine	
Category	Antikörper	
Product Description	The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule. In immunoelectrophoresis and double radial immunodiffusion, using various antiserum concentrations against normal human plasma and serum, a characteristic precipitin I	
Clonality	Monoclonal	
Clone Designation	[Polyclonal]	
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added. Reconstitute the lyophilized	
Source	Purified normal IgG isolated from pooled human serum. Freunds complete adjuvant is used in the first step of the immunization procedure.	

Aр	plication	Notes

Precipitation assays. In immunoelectrophoresis use 2  $\mu l$  or equivalent against 120  $\mu l$  antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10  $\mu l$  in a 3 mm diameter center well and 2  $\mu l$  serum samples (neat and serially d