

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Swine anti Human IgM (Fc specific), Clone: [Polyclonal], Porcine, Monoclonal NMB-SWAHU/IGM(FC)

Article Name	Swine anti Human IgM (Fc specific), Clone: [Polyclonal], Porcine,
	Monoclonal
Biozol Catalog Number	NMB-SWAHU/IGM(FC)
Supplier Catalog Number	SwAHu/IgM(Fc)
Alternative Catalog Number	NMB-SWAHU/IGM(FC)
Manufacturer	NordicMubio
Host	Porcine
Category	Antikörper
Product Description	The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal human serum a single precipitin line is obtained which shows
Clonality	Monoclonal
Clone Designation	[Polyclonal]
Buffer	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added. Reconstitute the lyophilized
Source	Highly purified normal IgM isolated from pooled human serum. Freunds complete adjuvant is used in the first step of the immunization procedure.

Formula	Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal swine serum. No foreign proteins added.
Application Notes	Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l in a 3 mm diameter center well and 2 $\mu$ l serum samples (neat and serially d