

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD20 Purified Azide Free, Clone: [MEM-97], Monoclonal EXB-10-217-C100

Article Name       Anti-Hu CD20 Purified Azide Free, Clone: [MEM-97], Monoclonal         Biozol Catalog Number       EXB-10-217-C100         Supplier Catalog Number       EXB-10-217-C100         Alternative Catalog Number       EXB-10-217-C100         Manufacturer       EXBIO         Category       Antikörper         Application       FC, IP         Species Reactivity       Bovine, Human         Immunogen       Raji human Burkitts lymphoma cell line         CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphorytetin expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on         Clonality       Monoclonal         Concentration       1 mg/ml         Clone Designation       [MEM-97]         Isotype       Mouse IgG1         Buffer       Phosphate buffered saline (PBS), pH 7.4         Storage       2°C to 8°C		
Supplier Catalog Number 10-217-C100  Alternative Catalog Number EXB-10-217-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IP  Species Reactivity Bovine, Human  Immunogen Raji human Burkitts lymphoma cell line  CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-97]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4	Article Name	Anti-Hu CD20 Purified Azide Free, Clone: [MEM-97], Monoclonal
Alternative Catalog Number   EXB-10-217-C100  Manufacturer   EXBIO  Category   Antikörper  Application   FC, IP  Species Reactivity   Bovine, Human  Immunogen   Raji human Burkitts lymphoma cell line  CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on  Clonality   Monoclonal  Concentration   I mg/ml  Clone Designation   [MEM-97]  Isotype   Mouse IgG1  Phosphate buffered saline (PBS), pH 7.4	Biozol Catalog Number	EXB-10-217-C100
Manufacturer  EXBIO  Category  Antikörper  Application  FC, IP  Species Reactivity  Bovine, Human  Immunogen  Raji human Burkitts lymphoma cell line  CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on  Clonality  Monoclonal  Concentration  I mg/ml  Clone Designation  [MEM-97]  Isotype  Mouse IgG1  Buffer  Phosphate buffered saline (PBS), pH 7.4	Supplier Catalog Number	10-217-C100
Category Antikörper  Application FC, IP  Species Reactivity Bovine, Human  Immunogen Raji human Burkitts lymphoma cell line  CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-97]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4	Alternative Catalog Number	EXB-10-217-C100
Application FC, IP  Species Reactivity Bovine, Human  Immunogen Raji human Burkitts lymphoma cell line  CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-97]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4	Manufacturer	EXBIO
Species Reactivity  Bovine, Human  Raji human Burkitts lymphoma cell line  CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [MEM-97]  Isotype  Mouse IgG1  Buffer  Phosphate buffered saline (PBS), pH 7.4	Category	Antikörper
Immunogen       Raji human Burkitts lymphoma cell line         CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on         Clonality       Monoclonal         Concentration       1 mg/ml         Clone Designation       [MEM-97]         Isotype       Mouse IgG1         Buffer       Phosphate buffered saline (PBS), pH 7.4	Application	FC, IP
CD20 is a cell surface 33-37 (depending on the degree of phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-97]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4	Species Reactivity	Bovine, Human
phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected on  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [MEM-97]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4	Immunogen	Raji human Burkitts lymphoma cell line
Concentration 1 mg/ml Clone Designation [MEM-97] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4	Product Description	phosphorylation) kDa non-glycosylated surface phosphoprotein expressed on mature and most malignant B cells, but not stem cells or plasma cells (low number of the CD20 has been also detected
Clone Designation [MEM-97]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4	Clonality	Monoclonal
Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4	Clone Designation	[MEM-97]
	Isotype	Mouse IgG1
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4
	Storage	2°C to 8°C

Target	CD20
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-5 μg/ml.