

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CD3 zeta Purified Azide Free, Clone: [H146-968], Monoclonal EXB-10-568-C100

Article Name Anti-CD3 zeta Purified Azide Free, Clone: [H146-968], Monoclonal Biozol Catalog Number EXB-10-568-C100 Alternative Catalog Number EXB-10-568-C100 Manufacturer EXBIO Category Antikörper Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4 Storage 2°C to 8°C		
Supplier Catalog Number 10-568-C100 Alternative Catalog Number EXB-10-568-C100 Manufacturer EXBIO Category Antikörper Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen Synthetic peptide corresponding to amino acids 151-164 of mouse CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Article Name	Anti-CD3 zeta Purified Azide Free, Clone: [H146-968], Monoclonal
Alternative Catalog Number EXB-10-568-C100 Manufacturer EXBIO Category Antikörper Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen Synthetic peptide corresponding to amino acids 151-164 of mouse CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Biozol Catalog Number	EXB-10-568-C100
Manufacturer EXBIO Category Antikörper Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen Synthetic peptide corresponding to amino acids 151-164 of mouse CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Supplier Catalog Number	10-568-C100
Category Antikörper Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen Synthetic peptide corresponding to amino acids 151-164 of mouse CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Alternative Catalog Number	EXB-10-568-C100
Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen Synthetic peptide corresponding to amino acids 151-164 of mouse CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Manufacturer	EXBIO
Species Reactivity Human, Mouse Synthetic peptide corresponding to amino acids 151-164 of mouse CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Category	Antikörper
Immunogen Synthetic peptide corresponding to amino acids 151-164 of mouse CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Application	FC, ICC, IP, WB
Product Description CD3 zeta. CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Species Reactivity	Human, Mouse
Product Description into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of t Clonality Monoclonal Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Immunogen	
Concentration 1 mg/ml Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Product Description	into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen
Clone Designation [H146-968] Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Clonality	Monoclonal
Isotype Armenian Hamster IgG2 kappa Buffer Phosphate buffered saline (PBS), pH 7.4	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4	Clone Designation	[H146-968]
	Isotype	Armenian Hamster IgG2 kappa
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4
	Storage	2°C to 8°C

Target	CD3 zeta
Antibody Type	Monoclonal Antibody
Application Notes	Immunocytochemistry: Recommended dilution: 4 μ g/ml. Western blotting: Recommended dilution: 1-2 μ g/ml, reducing conditions, positive control: MOLT-4 cells, murine splenocytes. Flow cytometry: Recommended dilution: 1-5 μ g/ml. Intracellular stainin