

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Cytokeratin 18 Purified, Clone: [DA-7], Monoclonal EXB-11-110-C025

Article Name Anti-Cytokeratin 18 Purified, Clone: [DA-7], Monoclonal Biozol Catalog Number EXB-11-110-C025 Supplier Catalog Number 11-110-C025 Alternative Catalog Number EXB-11-110-C025 Manufacturer EXBIO Category Antikörper Application IP, ELISA, IHC-P, ICC, WB Species Reactivity Human Immunogen Human breast carcinoma cell line PMC-42. Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The indivi
Supplier Catalog Number Alternative Catalog Number EXB-11-110-C025 Manufacturer EXBIO Category Antikörper Application IP, ELISA, IHC-P, ICC, WB Species Reactivity Human Immunogen Human breast carcinoma cell line PMC-42. Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
Alternative Catalog Number EXB-11-110-C025 Manufacturer EXBIO Category Antikörper Application IP, ELISA, IHC-P, ICC, WB Species Reactivity Human Immunogen Human breast carcinoma cell line PMC-42. Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
Manufacturer EXBIO Category Antikörper Application IP, ELISA, IHC-P, ICC, WB Species Reactivity Human Human breast carcinoma cell line PMC-42. Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
Category Application IP, ELISA, IHC-P, ICC, WB Species Reactivity Human Human breast carcinoma cell line PMC-42. Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
Application IP, ELISA, IHC-P, ICC, WB Species Reactivity Human Immunogen Human breast carcinoma cell line PMC-42. Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
Species Reactivity Human Human breast carcinoma cell line PMC-42. Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
Immunogen Human breast carcinoma cell line PMC-42. Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
Cytokeratins are a subfamily of intermediate filaments and are characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
characterized by remarkable biochemical diversity. They are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The
marrin.
Clonality Monoclonal
Concentration 1 mg/ml
Clone Designation [DA-7]
Isotype IgG1
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage 2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Western blotting: Recommended dilution: 1-2 $\mu\text{g/ml.Immunohistochemistry}$ (paraffin sections): Recommended dilution: 10 $\mu\text{g/ml}$, positive tissue: colon, skin. Immunocytochemistry: Positive control: MCF-7 human breast adenocarcinoma cell line, HeLa hum