

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-CDK1 Purified, Clone: [POH-1], Monoclonal EXB-11-115-C100

Anti-CDK1 Purified, Clone: [POH-1], Monoclonal
EXB-11-115-C100
11-115-C100
EXB-11-115-C100
EXBIO
Antikörper
ICC, IHC-P, IP, WB
Bovine, Human, Primate
Bacterially expressed full-length human Cdk1 protein
Cdk1 (cyclin-dependent kinase 1), also known as p34Cdc2 (cell division control protein kinase 2) depends on cyclin A and B and is triggered by a positive feedback loop at the end of G2 phase, which is the key event that initiates mitotic entry. Destr
Monoclonal
1 mg/ml
[POH-1]
Mouse IgG2a
Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
2°C to 8°C

Target	CDK1
Antibody Type	Monoclonal Antibody
Application Notes	Immunohistochemistry (paraffin sections): Recommended dilution: 10 μ g/ml, positive tissue: thymus.Western blotting: Recommended dilution: 1-2 μ g/ml.