

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD157 Purified, Clone: [SY11B5], Monoclonal EXB-11-136-C100

Article Name Anti-Hu CD157 Purified, Clone: [SY11B5], Monoclonal Biozol Catalog Number EXB-11-136-C100 Alternative Catalog Number EXB-11-136-C100 Alternative Catalog Number EXB-11-136-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human, Primate Immunogen Human CD157 CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C		
Supplier Catalog Number 11-136-C100 Alternative Catalog Number EXB-11-136-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human, Primate Immunogen Human CD157 CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD157 Purified, Clone: [SY11B5], Monoclonal
Alternative Catalog Number EXB-11-136-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human, Primate Immunogen Human CD157 CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-136-C100
Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human, Primate Immunogen Human CD157 CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-136-C100
Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human, Primate Immunogen Human CD157 CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-136-C100
Application FC, IHC-Fr, IP, WB Species Reactivity Human, Primate Immunogen Human CD157 CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Human, Primate Human CD157 CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen Human CD157 CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC, IHC-Fr, IP, WB
CD157 (cADPr hydrolase 2) is a GPI-anchored ectoenzyme possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Human, Primate
Product Description possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular dendriti Clonality Monoclonal Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	Human CD157
Concentration 1 mg/ml Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	possessing ADP-ribosyl cyclase and cyclic ADP-ribose hydrolase activity. It uses NAD and cADP-ribose as substrates. CD157 is expressed on granulocytes, monocytes, macrophages, follicular
Clone Designation [SY11B5] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[SY11B5]
	Isotype	Mouse IgG1 kappa
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
	Storage	2°C to 8°C

Target	CD157
Antibody Type	Monoclonal Antibody
Application Notes	Western blotting: Non-reducing conditions. Immunohistochemistry: Recommended dilution: 5-10 μ g/ml. Flow cytometry: Recommended dilution: 1-12 μ g/ml