

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD49c Purified, Clone: [ASC-1], Monoclonal EXB-11-137-C100

Article Name Anti-Hu CD49c Purified, Clone: [ASC-1], Monoclonal Biozol Catalog Number EXB-11-137-C100 Supplier Catalog Number 11-137-C100 Alternative Catalog Number EXB-11-137-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human Immunogen Human SSC-9 cell line (squamous cell carcinoma) CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration 1 mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide		
Supplier Catalog Number Alternative Catalog Number EXB-11-137-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human Immunogen Human SSC-9 cell line (squamous cell carcinoma) CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration I mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD49c Purified, Clone: [ASC-1], Monoclonal
Alternative Catalog Number EXB-11-137-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human Immunogen Human SSC-9 cell line (squamous cell carcinoma) CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration I mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-137-C100
Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human Human SSC-9 cell line (squamous cell carcinoma) CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration I mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-137-C100
Category Antikörper Application FC, IHC-Fr, IP, WB Species Reactivity Human Immunogen Human SSC-9 cell line (squamous cell carcinoma) CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration 1 mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-137-C100
Application FC, IHC-Fr, IP, WB Species Reactivity Human Human SSC-9 cell line (squamous cell carcinoma) CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration I mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Immunogen Human SSC-9 cell line (squamous cell carcinoma) CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration I mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen Human SSC-9 cell line (squamous cell carcinoma) CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration I mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC, IHC-Fr, IP, WB
Product Description CD49c / Integrin alpha 3 is a type I transmembrane glycoprotein proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration 1 mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Human
Product Description proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-3 complex, an adhesion receptor for extracellular matrix Clonality Monoclonal Concentration 1 mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	Human SSC-9 cell line (squamous cell carcinoma)
Concentration 1 mg/ml Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	proteolytically cleaved into two disulfide linked chains. It noncovalently associates with CD29 (integrin beta 1) to form the VLA-
Clone Designation [ASC-1] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[ASC-1]
	Isotype	Mouse IgG1 kappa
	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage 2°C to 8°C	Storage	2°C to 8°C

Target	CD49c
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-4 μg/ml.