

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD89 Purified, Clone: [A59], Monoclonal EXB-11-139-C025

Article Name Anti-Hu CD89 Purified, Clone: [A59], Monoclonal Biozol Catalog Number EXB-11-139-C025  Supplier Catalog Number 11-139-C025  Alternative Catalog Number EXB-11-139-C025  Manufacturer EXBIO  Category Antikörper  Application FC  Species Reactivity Primate, Human  Immunogen Ag8.653 myeloma cells  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality Monoclonal  Concentration I mg/ml  Clone Designation [A59]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide		
Supplier Catalog Number  Alternative Catalog Number  EXB-11-139-C025  Manufacturer  EXBIO  Category  Antikörper  Application  FC  Species Reactivity  Primate, Human  Immunogen  Ag8.653 myeloma cells  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality  Monoclonal  Concentration  I mg/ml  Clone Designation  [A59]  Isotype  IgG1, kappa  Buffer  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD89 Purified, Clone: [A59], Monoclonal
Alternative Catalog Number EXB-11-139-C025  Manufacturer EXBIO  Category Antikörper  Application FC  Species Reactivity Primate, Human  Immunogen Ag8.653 myeloma cells  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [A59]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-139-C025
Manufacturer  EXBIO  Category  Antikörper  Application  FC  Species Reactivity  Primate, Human  Ag8.653 myeloma cells  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [A59]  Isotype  IgG1, kappa  Buffer  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-139-C025
Category Antikörper  Application FC  Species Reactivity Primate, Human  Immunogen Ag8.653 myeloma cells  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [A59]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-139-C025
Application FC  Species Reactivity Primate, Human  Immunogen Ag8.653 myeloma cells  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [A59]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity  Immunogen  Ag8.653 myeloma cells  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality  Monoclonal  Concentration  I mg/ml  Clone Designation  [A59]  Isotype  IgG1, kappa  Buffer  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen  Ag8.653 myeloma cells  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality  Monoclonal  Concentration  I mg/ml  Clone Designation  [A59]  Isotype  IgG1, kappa  Buffer  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC
Product Description  CD89 (Fc-alpha-R) is a type I transmembrane glycoprotein serving as a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [A59]  Isotype  IgG1, kappa  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Primate, Human
Product Description  a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with FcR gamma chain homodimers is needed. CD89  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [A59]  Isotype  IgG1, kappa  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	Ag8.653 myeloma cells
Concentration 1 mg/ml  Clone Designation [A59]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	a receptor for IgA. Soluble CD89 is detectable in serum and retains its IgA binding capacity. For signal transduction the association with
Clone Designation [A59]  Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype IgG1, kappa  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[A59]
	Isotype	IgG1, kappa
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
2 0 0 0 0	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-4 μg/ml.