

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD137 Purified, Clone: [4B4-1], Monoclonal EXB-11-143-C025

Article Name Anti-Hu CD137 Purified, Clone: [4B4-1], Monoclonal Biozol Catalog Number EXB-11-143-C025 Supplier Catalog Number 11-143-C025 Alternative Catalog Number EXB-11-143-C025 Manufacturer EXBIO Category Antikörper Application FC, ELISA, IP Species Reactivity Primate, Human Immunogen Recombinant human CD137 ectodomain CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C		
Supplier Catalog Number 11-143-C025 Alternative Catalog Number EXB-11-143-C025 Manufacturer EXBIO Category Antikörper Application FC, ELISA, IP Species Reactivity Primate, Human Immunogen Recombinant human CD137 ectodomain CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD137 Purified, Clone: [4B4-1], Monoclonal
Alternative Catalog Number EXB-11-143-C025 Manufacturer EXBIO Category Antikörper Application FC, ELISA, IP Species Reactivity Primate, Human Immunogen Recombinant human CD137 ectodomain CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-143-C025
Manufacturer EXBIO Category Antikörper Application FC, ELISA, IP Species Reactivity Primate, Human Recombinant human CD137 ectodomain CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-143-C025
Category Antikörper Application FC, ELISA, IP Species Reactivity Primate, Human Immunogen Recombinant human CD137 ectodomain CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-143-C025
Application FC, ELISA, IP Species Reactivity Primate, Human Immunogen Recombinant human CD137 ectodomain CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Primate, Human Recombinant human CD137 ectodomain CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
ImmunogenRecombinant human CD137 ectodomainProduct DescriptionCD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitorClonalityMonoclonalConcentration1 mg/mlClone Designation[4B4-1]IsotypeIgG1, kappaBufferPhosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC, ELISA, IP
Product Description CD137, also known as TNFRSF9 or 4-1BB, is an inducible costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Primate, Human
Product Description costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages, mature B cells, hematopoietic stem cells, and myeloid progenitor Clonality Monoclonal Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	Recombinant human CD137 ectodomain
Concentration 1 mg/ml Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	costimulatory molecule expressed mainly on activated T cells. Its ligand, known as 4-1BBL, is expressed on activated macrophages,
Clone Designation [4B4-1] Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype IgG1, kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[4B4-1]
Thosphate barrered saline (1 55), pri 7.4, 13 mm socialii dzide	Isotype	IgG1, kappa
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-4 µg/ml