

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD30 Purified, Clone: [Ber-H8], Monoclonal EXB-11-153-C100

Article NameAnti-Hu CD30 Purified, Clone: [Ber-H8], MonoclonalBiozol Catalog NumberEXB-11-153-C100Supplier Catalog Number11-153-C100Alternative Catalog NumberEXB-11-153-C100ManufacturerEXBIOCategoryAntikörperApplicationFC, IHC-Fr, IHC-PSpecies ReactivityHumanProduct DescriptionCD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Kin. The ligand for CD30 is CD30L (CD153). The bindiClonalityMonoclonalConcentration1 mg/mlClone Designation[Ber-H8]IsotypeMouse IgG1 kappaBufferPhosphate buffered saline (PBS), pH 7.4, 15 mM sodium azideStorage2°C to 8°CTargetCD30		
Supplier Catalog Number 11-153-C100 Alternative Catalog Number EXB-11-153-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IHC-P Species Reactivity Human CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Kin. The ligand for CD30 is CD30L (CD153). The bindi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Article Name	Anti-Hu CD30 Purified, Clone: [Ber-H8], Monoclonal
Alternative Catalog Number EXB-11-153-C100 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IHC-P Species Reactivity Human CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki-1. The ligand for CD30 is CD30L (CD153). The bindi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Biozol Catalog Number	EXB-11-153-C100
Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr, IHC-P Species Reactivity Human CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki- 1. The ligand for CD30 is CD30L (CD153). The bindi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Supplier Catalog Number	11-153-C100
Category Antikörper Application FC, IHC-Fr, IHC-P Species Reactivity Human CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki-1. The ligand for CD30 is CD30L (CD153). The bindi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Alternative Catalog Number	EXB-11-153-C100
Application FC, IHC-Fr, IHC-P Species Reactivity Human CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki-1. The ligand for CD30 is CD30L (CD153). The bindi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Manufacturer	EXBIO
Species Reactivity Human CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki-1. The ligand for CD30 is CD30L (CD153). The bindi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Category	Antikörper
CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki- 1. The ligand for CD30 is CD30L (CD153). The bindi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Application	FC, IHC-Fr, IHC-P
Product Description superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki- 1. The ligand for CD30 is CD30L (CD153). The bindi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Species Reactivity	Human
Concentration 1 mg/ml Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Product Description	superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki-
Clone Designation [Ber-H8] Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Clonality	Monoclonal
Isotype Mouse IgG1 kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C	Clone Designation	[Ber-H8]
Storage 2°C to 8°C	Isotype	Mouse IgG1 kappa
	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Target CD30	Storage	2°C to 8°C
	Target	CD30

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-4 μg/ml. Immunohistochemistry: Recommended dilution: 5-10 μg/ml.