

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Cyclin D1 Purified, Clone: [DCS-6], Monoclonal EXB-11-166-C100

Anti-Cyclin D1 Purified, Clone: [DCS-6], Monoclonal
EXB-11-166-C100
11-166-C100
EXB-11-166-C100
EXBIO
Antikörper
FC, ICC, IHC-Fr, IHC-P, IP, WB
Human, Mouse, Primate, Rat
recombinant human cyclin D1 (amino acids 1-295)
Cyclin D1 (PRAD1, Bcl-1) is a cytoplasmic and nuclear protein, which is synthesized during G1 phase and assembles with either cyclin-dependent kinase 4 (CDK4) or CDK6 in response to growth factor stimulation. D-type cyclin-CDK complexes act to inacti
Monoclonal
1 mg/ml
[DCS-6]
Mouse IgG2a
Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
2°C to 8°C

Target	Cyclin D1
Antibody Type	Monoclonal Antibody
Application Notes	Immunohistochemistry: Recommended dilution: 5-10 μ g/ml. Flow cytometry: Recommended dilution: 3-12 μ g/ml. Intracellular staining.