

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD22 Purified, Clone: [S-HCL-1], Monoclonal EXB-11-175-C025

Anti-Hu CD22 Purified, Clone: [S-HCL-1], Monoclonal
EXB-11-175-C025
11-175-C025
EXB-11-175-C025
EXBIO
Antikörper
FC
Human
Whole hairy cell leukemia cells and membrane preparation
CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the
Monoclonal
1 mg/ml
[S-HCL-1]
IgG2b, kappa
Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Extracellular and intracellular staining, recommended dilution: 1-4 μ g/ml.