

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu beta2-Microglobulin Purified, Clone: [B2M-01], Monoclonal EXB-11-237-C100

Anti-Hu beta2-Microglobulin Purified, Clone: [B2M-01], Monoclonal
EXB-11-237-C100
11-237-C100
EXB-11-237-C100
EXBIO
Antikörper
ELISA, FC, ICC, IHC-P, IP, RIA, WB
Human
Purified human beta2-microglobulin
Beta2-microglobulin non-covalently associates with the 44 kDa alpha chain to forms the HLA Class I antigen complex. Human beta2-microglobulin associated with HLA Class I antigens is expressed on many types of cells including lymphocytes, thymocytes,
Monoclonal
1 mg/ml
[B2M-01]
Mouse IgG2a
Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
2°C to 8°C

Target	beta2-Microglobulin
Antibody Type	Monoclonal Antibody
Application Notes	Western blotting: Recommended dilution: 2-4 µg/ml, positive control: U937 human histiocytic lymphoma cell line, RAJI human Burkitt lymphoma cell line, negative control: EL4 mouse lymphoblastic lymphoma cell line, both reducing and non-reducing conditions