

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD22 Purified, Clone: [MEM-01], Monoclonal EXB-11-265-C100

Anti-Hu CD22 Purified, Clone: [MEM-01], Monoclonal
EXB-11-265-C100
11-265-C100
EXB-11-265-C100
EXBIO
Antikörper
FC, IP, WB
Human, Primate
Raji Burkitts lymphoma cell line
CD22, also known as Siglec-2 (sialic acid-binding immunoglobulin-like lectin-2) is a transmembrane glycoprotein binding alpha2,6-linked sialic acid-bearing ligands. Intracellular domain of CD22 recruits protein tyrosine phosphatase SHP-1 through the
Monoclonal
1 mg/ml
[MEM-01]
Mouse IgG1
Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
2°C to 8°C

Target	CD22
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 2-4 μ g/ml. Western blotting: Recommended dilution: 2-4 μ g/ml, 120 min on vertical incubator, positive control: RAJI human lymphoma cell line. The antibody MEM-01 stains only the zone corresponding to 140 kDa, but