

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD21 Purified, Clone: [LT21], Monoclonal EXB-11-306-C025

Article Name Anti-Hu CD21 Purified, Clone: [LT21], Monoclonal Biozol Catalog Number EXB-11-306-C025 Supplier Catalog Number 11-306-C025 Alternative Catalog Number EXB-11-306-C025 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr Species Reactivity Canine, Porcine, Human, Bovine Immunogen IM9 human B-lymphoblastoid cell line CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C		
Supplier Catalog Number 11-306-C025 Alternative Catalog Number EXB-11-306-C025 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr Species Reactivity Canine, Porcine, Human, Bovine Immunogen IM9 human B-lymphoblastoid cell line CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration Img/ml Clone Designation [LT21] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD21 Purified, Clone: [LT21], Monoclonal
Alternative Catalog Number EXB-11-306-C025 Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr Species Reactivity Canine, Porcine, Human, Bovine Immunogen IM9 human B-lymphoblastoid cell line CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-306-C025
Manufacturer EXBIO Category Antikörper Application FC, IHC-Fr Species Reactivity Canine, Porcine, Human, Bovine Immunogen IM9 human B-lymphoblastoid cell line CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-306-C025
Category Antikörper Application FC, IHC-Fr Species Reactivity Canine, Porcine, Human, Bovine Immunogen IM9 human B-lymphoblastoid cell line CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-306-C025
Application FC, IHC-Fr Species Reactivity Canine, Porcine, Human, Bovine Immunogen IM9 human B-lymphoblastoid cell line CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Canine, Porcine, Human, Bovine IM9 human B-lymphoblastoid cell line CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen IM9 human B-lymphoblastoid cell line CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC, IHC-Fr
CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Canine, Porcine, Human, Bovine
fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, s Clonality Monoclonal Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	IM9 human B-lymphoblastoid cell line
Concentration 1 mg/ml Clone Designation [LT21] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of
Clone Designation [LT21] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[LT21]
	Isotype	lgG1
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 2 µg/ml.