

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD41a Purified, Clone: [HIP8], Monoclonal EXB-11-417-C100

Article Name Anti-Hu CD41a Purified, Clone: [HIP8], Monoclonal  Biozol Catalog Number EXB-11-417-C100  Supplier Catalog Number 11-417-C100  Alternative Catalog Number EXB-11-417-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IHC-Fr  Species Reactivity Human, Primate  CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality Monoclonal  Concentration Img/ml  Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C  Target		
Supplier Catalog Number 11-417-C100  Alternative Catalog Number EXB-11-417-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IHC-Fr  Species Reactivity Human, Primate  CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Article Name	Anti-Hu CD41a Purified, Clone: [HIP8], Monoclonal
Alternative Catalog Number EXB-11-417-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IHC-Fr  Species Reactivity Human, Primate  CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Biozol Catalog Number	EXB-11-417-C100
Manufacturer  EXBIO  Category  Antikörper  Application  FC, IHC-Fr  Species Reactivity  Human, Primate  CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [HIP8]  Isotype  Mouse IgG1  Buffer  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage  2°C to 8°C	Supplier Catalog Number	11-417-C100
Category Antikörper  Application FC, IHC-Fr  Species Reactivity Human, Primate  CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Alternative Catalog Number	EXB-11-417-C100
Application FC, IHC-Fr  Species Reactivity Human, Primate  CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Manufacturer	EXBIO
Species Reactivity  Human, Primate  CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [HIP8]  Isotype  Mouse IgG1  Buffer  Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage  2°C to 8°C	Category	Antikörper
CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Application	FC, IHC-Fr
RDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a v  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Species Reactivity	Human, Primate
Concentration 1 mg/ml  Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Product Description	kDa a, alpha and 23 kDa b, beta) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor.
Clone Designation [HIP8]  Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Clonality	Monoclonal
Isotype Mouse IgG1  Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  Storage 2°C to 8°C	Clone Designation	[HIP8]
Storage 2°C to 8°C	Isotype	Mouse IgG1
	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Target CD41a	Storage	2°C to 8°C
	Target	CD41a

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-4 µg/ml