

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-Clusterin Purified, Clone: [Hs-3], Monoclonal EXB-11-421-C100

|                            |   |
|----------------------------|---|
| Article Name               | Anti-Clusterin Purified, Clone: [Hs-3], Monoclonal  |
| Biozol Catalog Number      | EXB-11-421-C100   |
| Supplier Catalog Number    | 11-421-C100   |
| Alternative Catalog Number | EXB-11-421-C100   |
| Manufacturer               | EXBIO   |
| Category                   | Antikörper  |
| Application                | ELISA, ICC, IHC-P, WB   |
| Species Reactivity         | Human   |
| Immunogen                  | Freshly ejaculated human sperms were washed in PBS and extracted in 3% acetic acid, 10% glycerol, 30 mM benzaminidine. The acid extract was dialyzed against 0.2% acetic acid and subsequently used for immunization.   |
| Product Description        | Clusterin (APO J, SGP-2, TRPM-2, SP-40, pADHC-9, CLJ, T64, GP III, XIP8) is a 75-80 kD disulfide-linked heterodimeric protein containing about 30% of N-linked carbohydrate rich in sialic acid but truncated forms targeted to the nucleus have also been... |
| Clonality                  | Monoclonal  |
| Concentration              | 1 mg/ml   |
| Clone Designation          | [Hs-3]  |
| Isotype                    | Mouse IgG1  |
| Buffer                     | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |

|                   |   |
|-------------------|---|
| Storage           | 2°C to 8°C  |
| Target            | Clusterin   |
| Antibody Type     | Monoclonal Antibody   |
| Application Notes | Immunohistochemistry (paraffin sections): Recommended dilution: 10 µg/ml, positive tissue: Sertoli cells. Western blotting: Recommended dilution: 1 µg/ml, positive material: seminal plasma. |