

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD86 Purified, Clone: [BU63], Monoclonal EXB-11-531-C025

Article Name Anti-Hu CD86 Purified, Clone: [BU63], Monoclonal Biozol Catalog Number EXB-11-531-C025 Supplier Catalog Number 11-531-C025 Alternative Catalog Number EXB-11-531-C025 Manufacturer EXBIO Category Antikörper Application WB, FC, IHC-Fr, IP Species Reactivity Human Immunogen B-lymphoblastoid cell line ARH 77 CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C		
Supplier Catalog Number 11-531-C025 Alternative Catalog Number EXB-11-531-C025 Manufacturer EXBIO Category Antikörper Application WB, FC, IHC-Fr, IP Species Reactivity Human Immunogen B-lymphoblastoid cell line ARH 77 CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD86 Purified, Clone: [BU63], Monoclonal
Alternative Catalog Number EXB-11-531-C025 Manufacturer EXBIO Category Antikörper Application WB, FC, IHC-Fr, IP Species Reactivity Human Immunogen B-lymphoblastoid cell line ARH 77 CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration I mg/ml Clone Designation [BU63] Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-531-C025
Manufacturer EXBIO Category Antikörper Application WB, FC, IHC-Fr, IP Species Reactivity Human Immunogen B-lymphoblastoid cell line ARH 77 CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration I mg/ml Clone Designation [BU63] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-531-C025
Category Application WB, FC, IHC-Fr, IP Species Reactivity Human B-lymphoblastoid cell line ARH 77 CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-531-C025
Application WB, FC, IHC-Fr, IP Species Reactivity Human Immunogen B-lymphoblastoid cell line ARH 77 CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Human B-lymphoblastoid cell line ARH 77 CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen B-lymphoblastoid cell line ARH 77 CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	WB, FC, IHC-Fr, IP
Product Description CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Human
Product Description costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, Clonality Monoclonal Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	B-lymphoblastoid cell line ARH 77
Concentration 1 mg/ml Clone Designation [BU63] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional
Clone Designation [BU63] Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[BU63]
	Isotype	lgG1
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
	Storage	2°C to 8°C

Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 5 μg/ml.