

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-PSMA Purified, Clone: [GCP-04], Monoclonal EXB-11-532-C025

| Article Name | Anti-PSMA Purified, Clone: [GCP-04], Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | EXB-11-532-C025 |
| Supplier Catalog Number | 11-532-C025 |
| Alternative Catalog Number | EXB-11-532-C025 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | WB, IHC-P, ICC |
| Species Reactivity | Mouse, Porcine, Human, Rat |
| Immunogen | Recombinant fragment of human GCPII (amino acids 44-750) produced in S2 cells |
| Product Description | Glutamate carboxypeptidase II (GCPII), also known as N-acetyl-alphalinked acidic dipeptidase I (NAALADase I), folate hydrolase (FOLH1), and prostate-specific membrane antigen (PSMA), is an approximately 95-110 kDa type II transmembrane glycoprotein |
| Clonality | Monoclonal |
| Concentration | 1 mg/ml |
| Clone Designation | [GCP-04] |
| Isotype | IgG1 |
| Buffer | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage | 2°C to 8°C |

| Antibody Type | Monoclonal Antibody |
|-------------------|---|
| Application Notes | Western blotting: Recommended dilution: 1 μ g/ml, positive control: LNCaP cell line. Sample preparation: Resuspend approx. 50 mil. cells in 1 ml cold lysis buffer (1% NP-40). Incubate 30 min on ice. Mix lysate with non-reducing/reducing Laemmli SDS-PAGE s |