

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu IgG (Fc) Purified, Clone: [EM-07], Monoclonal EXB-11-536-C100

Article Name Anti-Hu IgG (Fc) Purified, Clone: [EM-07], Monoclonal Biozol Catalog Number EXB-11-536-C100 Supplier Catalog Number 11-536-C100 Alternative Catalog Number EXB-11-536-C100 Manufacturer EXBIO Category Antikörper Application ELISA, FC, ICC, WB Species Reactivity Human Immunogen Fusion protein of human IgG Fc fragment. Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide		
Supplier Catalog Number Alternative Catalog Number EXB-11-536-C100 Manufacturer EXBIO Category Antikörper Application ELISA, FC, ICC, WB Species Reactivity Human Immunogen Fusion protein of human IgG Fc fragment. Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu IgG (Fc) Purified, Clone: [EM-07], Monoclonal
Alternative Catalog Number EXB-11-536-C100 Manufacturer EXBIO Category Antikörper Application ELISA, FC, ICC, WB Species Reactivity Human Immunogen Fusion protein of human IgG Fc fragment. Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration I mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-536-C100
Manufacturer EXBIO Category Antikörper Application ELISA, FC, ICC, WB Species Reactivity Human Fusion protein of human IgG Fc fragment. Immunogen Fusion protein of human IgG Fc fragment. Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-536-C100
Category Antikörper Application ELISA, FC, ICC, WB Species Reactivity Human Immunogen Fusion protein of human IgG Fc fragment. Product Description Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-536-C100
Application ELISA, FC, ICC, WB Species Reactivity Human Fusion protein of human IgG Fc fragment. Immunogen Fusion protein of human IgG Fc fragment. Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Immunogen Fusion protein of human IgG Fc fragment. Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration I mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen Fusion protein of human IgG Fc fragment. Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration I mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	ELISA, FC, ICC, WB
Product Description Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Human
Product Description major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the Clonality Monoclonal Concentration 1 mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	Fusion protein of human IgG Fc fragment.
Concentration 1 mg/ml Clone Designation [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is
Clone Designation [EM-07] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[EM-07]
	Isotype	Mouse IgG1
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Z C to 6 C	Storage	2°C to 8°C

Target	IgG (Fc)
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1-4 μg/ml.