

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Hu CD69 Purified, Clone: [FN50], Monoclonal EXB-11-552-C100

| Article Name               | Anti III. CDCO Durified Clane. [FNEO] Marantanal   |
|----------------------------|--|
| Article Name               | Anti-Hu CD69 Purified, Clone: [FN50], Monoclonal   |
| Biozol Catalog Number      | EXB-11-552-C100  |
| Supplier Catalog Number    | 11-552-C100  |
| Alternative Catalog Number | EXB-11-552-C100  |
| Manufacturer               | EXBIO  |
| Category                   | Antikörper   |
| Application                | FC, IHC-Fr   |
| Species Reactivity         | Human  |
| Immunogen                  | anti-µ-stimulated human B lymphocytes  |
| Product Description        | CD69 (C-type lectin domain family 2 C, CLEC2C, also known as AIM) is one of the earliest inducible cell surface molecules acquired during leukocyte activation. This glycoprotein serves as a lectin-type receptor in lymphocytes, NK cells and platelets, |
| Clonality                  | Monoclonal   |
| Concentration              | 1 mg/ml  |
| Clone Designation          | [FN50]   |
| Isotype                    | Mouse IgG1   |
| Buffer                     | Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide  |
| Storage                    | 2°C to 8°C   |

| Target             | CD69  |
|--------------------|---|
| Antibody Type      | Monoclonal Antibody   |
| Application Dilute | FUNCWhen stimulating peripheral blood T cells, FN50 antibody together with TPA synergistically increases cell volume and RNA/DNA synthesis. Addition of FN50 to stimulated peripheral blood B cells has similar, but weaker effect. |
| Application Notes  | Flow cytometry: recommended dilution: 1-4 µg/ml.  |