

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD162 Purified, Clone: [TC2], Monoclonal EXB-11-603-C100

Article Name Anti-Hu CD162 Purified, Clone: [TC2], Monoclonal Biozol Catalog Number EXB-11-603-C100 Supplier Catalog Number 11-603-C100 Alternative Catalog Number EXB-11-603-C100 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen Human thymocytes CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C		
Supplier Catalog Number 11-603-C100 Alternative Catalog Number EXB-11-603-C100 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen Human thymocytes CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Hu CD162 Purified, Clone: [TC2], Monoclonal
Alternative Catalog Number EXB-11-603-C100 Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Immunogen Human thymocytes CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration I mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-603-C100
Manufacturer EXBIO Category Antikörper Application FC Species Reactivity Human Human thymocytes CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration I mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-603-C100
Category Antikörper Application FC Species Reactivity Human Immunogen Human thymocytes CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-603-C100
Application FC Species Reactivity Human Immunogen Human thymocytes CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Immunogen Human thymocytes CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen Human thymocytes CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC
Product Description CD162 (P-selectin glycoprotein ligand-1, PSGL-1) is a sialomucin constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Human
Product Description constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves as a ligand for P- E- and L-selectin, with the hi Clonality Monoclonal Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	Human thymocytes
Concentration 1 mg/ml Clone Designation [TC2] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	constitutively expressed as a disulfide-linked homodimer of two 120 kDa subunits on the surface of circulating leukocytes. CD162 serves
Clone Designation [TC2] Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype Mouse IgG1 Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[TC2]
	Isotype	Mouse IgG1
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
	Storage	2°C to 8°C

Target	CD162
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1.5 - 2.5 $\mu g/ml$, positive control: human peripheral blood.