

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Anti-Ms CD11a Purified, Clone: [M17/4], Monoclonal EXB-11-622-C100

Anti-Ms CD11a Purified, Clone: [M17/4], Monoclonal Biozol Catalog Number EXB-11-622-C100  Supplier Catalog Number 11-622-C100  Alternative Catalog Number EXB-11-622-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IHC-Fr, IP  Species Reactivity Mouse  Immunogen C57BL/6 mouse splenic secondary cytotoxic T lymphocytes  CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphal.beta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [M17/4]  Isotype Rat IgG2a		
Supplier Catalog Number 11-622-C100  Alternative Catalog Number EXB-11-622-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IHC-Fr, IP  Species Reactivity Mouse  Immunogen C57BL/6 mouse splenic secondary cytotoxic T lymphocytes  CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [M17/4]	Article Name	Anti-Ms CD11a Purified, Clone: [M17/4], Monoclonal
Alternative Catalog Number EXB-11-622-C100  Manufacturer EXBIO  Category Antikörper  Application FC, IHC-Fr, IP  Species Reactivity Mouse  Immunogen C57BL/6 mouse splenic secondary cytotoxic T lymphocytes  CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [M17/4]	Biozol Catalog Number	EXB-11-622-C100
Manufacturer  EXBIO  Category  Antikörper  Application  FC, IHC-Fr, IP  Species Reactivity  Mouse  Immunogen  C57BL/6 mouse splenic secondary cytotoxic T lymphocytes  CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity  Clonality  Monoclonal  Concentration  I mg/ml  Clone Designation  [M17/4]	Supplier Catalog Number	11-622-C100
Category Antikörper  Application FC, IHC-Fr, IP  Species Reactivity Mouse  Immunogen C57BL/6 mouse splenic secondary cytotoxic T lymphocytes  CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [M17/4]	Alternative Catalog Number	EXB-11-622-C100
Application FC, IHC-Fr, IP  Species Reactivity Mouse  Immunogen C57BL/6 mouse splenic secondary cytotoxic T lymphocytes  CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity  Clonality Monoclonal  Concentration 1 mg/ml  Clone Designation [M17/4]	Manufacturer	EXBIO
Species Reactivity       Mouse         Immunogen       C57BL/6 mouse splenic secondary cytotoxic T lymphocytes         Product Description       CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity         Clonality       Monoclonal         Concentration       1 mg/ml         Clone Designation       [M17/4]	Category	Antikörper
Immunogen       C57BL/6 mouse splenic secondary cytotoxic T lymphocytes         Product Description       CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity         Clonality       Monoclonal         Concentration       1 mg/ml         Clone Designation       [M17/4]	Application	FC, IHC-Fr, IP
Product Description  CD11a (LFA-1 alpha) together with CD18 constitute leukocyte function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [M17/4]	Species Reactivity	Mouse
Product Description  function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin.  CD11a is implicated in activation of LFA-1 complex. LFA-1 is expressed on the plasma membrane of leukocytes in a low-affinity  Clonality  Monoclonal  Concentration  1 mg/ml  Clone Designation  [M17/4]	Immunogen	C57BL/6 mouse splenic secondary cytotoxic T lymphocytes
Concentration 1 mg/ml  Clone Designation [M17/4]	Product Description	function-associated antigen 1 (LFA-1), the alphaLbeta2 integrin. CD11a is implicated in activation of LFA-1 complex. LFA-1 is
Clone Designation [M17/4]	Clonality	Monoclonal
	Concentration	1 mg/ml
Isotype Rat IqG2a	Clone Designation	[M17/4]
The state of the s	Isotype	Rat IgG2a
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage 2°C to 8°C	Storage	2°C to 8°C

Target	CD11a
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1 $\mu$ g/ml. Immunohistochemistry (frozen sections): Positive tissue: murine spleen or thymus, acetone fixation.