

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Ms CD44 Purified, Clone: [IM7], Monoclonal EXB-11-624-C100

Article Name Anti-Ms CD44 Purified, Clone: [IM7], Monoclonal Biozol Catalog Number EXB-11-624-C100 Supplier Catalog Number 11-624-C100 Alternative Catalog Number EXB-11-624-C100 Manufacturer EXBIO Category Antikörper Application FC, ICC, IHC-Fr, IHC-P, IP, WB Species Reactivity Canine, Equine, Feline, Human, Mouse Immunogen Dexamethasone-induced cells of the SJL mouse spontaneous myeloid leukemia M1 CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide Storage 2°C to 8°C		
Supplier Catalog Number 11-624-C100 Alternative Catalog Number EXB-11-624-C100 Manufacturer EXBIO Category Antikörper Application FC, ICC, IHC-Fr, IHC-P, IP, WB Species Reactivity Canine, Equine, Feline, Human, Mouse Immunogen Dexamethasone-induced cells of the SJL mouse spontaneous myeloid leukemia M1 CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Article Name	Anti-Ms CD44 Purified, Clone: [IM7], Monoclonal
Alternative Catalog Number EXB-11-624-C100 Manufacturer EXBIO Category Antikörper Application FC, ICC, IHC-Fr, IHC-P, IP, WB Species Reactivity Canine, Equine, Feline, Human, Mouse Immunogen Dexamethasone-induced cells of the SJL mouse spontaneous myeloid leukemia M1 CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Biozol Catalog Number	EXB-11-624-C100
Manufacturer EXBIO Category Antikörper Application FC, ICC, IHC-Fr, IHC-P, IP, WB Species Reactivity Canine, Equine, Feline, Human, Mouse Immunogen Dexamethasone-induced cells of the SJL mouse spontaneous myeloid leukemia M1 CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration I mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Supplier Catalog Number	11-624-C100
Category Antikörper Application FC, ICC, IHC-Fr, IHC-P, IP, WB Species Reactivity Canine, Equine, Feline, Human, Mouse Immunogen Dexamethasone-induced cells of the SJL mouse spontaneous myeloid leukemia M1 CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Alternative Catalog Number	EXB-11-624-C100
Application FC, ICC, IHC-Fr, IHC-P, IP, WB Species Reactivity Canine, Equine, Feline, Human, Mouse Immunogen Dexamethasone-induced cells of the SJL mouse spontaneous myeloid leukemia M1 CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Manufacturer	EXBIO
Species Reactivity Canine, Equine, Feline, Human, Mouse Dexamethasone-induced cells of the SJL mouse spontaneous myeloid leukemia M1 CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Category	Antikörper
Immunogen Dexamethasone-induced cells of the SJL mouse spontaneous myeloid leukemia M1 Product Description CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Application	FC, ICC, IHC-Fr, IHC-P, IP, WB
Product Description CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Species Reactivity	Canine, Equine, Feline, Human, Mouse
Product Description most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculat Clonality Monoclonal Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Immunogen	
Concentration 1 mg/ml Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Product Description	most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it
Clone Designation [IM7] Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clonality	Monoclonal
Isotype Rat IgG2b kappa Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Concentration	1 mg/ml
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Clone Designation	[IM7]
	Isotype	Rat IgG2b kappa
Storage 2°C to 8°C	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
	Storage	2°C to 8°C

Target	CD44
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution: 1 μ g/ml, positive control: murine peripheral blood. Immunohistochemistry (paraffin sections): Positive tissue: murine spleen. Western blotting: Non-reducing conditions.