

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-ZAP70 Purified, Clone: [1E7.2], Monoclonal EXB-11-665-C100

Article NameAnti-ZAP70 Purified, Clone: [1E7.2], MonoclonalBiozol Catalog NumberEXB-11-665-C100Supplier Catalog Number11-665-C100Alternative Catalog NumberEXB-11-665-C100ManufacturerEXBIOCategoryAntikörperApplicationFC, ICC, IP, WBSpecies ReactivityHuman, MouseImmunogenA KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP70Product DescriptionThe ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase prClonalityMonoclonalConcentration1 mg/mlClone Designation[1E7.2]IsotypeMouse IgG1 kappaBufferPhosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide		
Supplier Catalog Number 11-665-C100 Alternative Catalog Number EXB-11-665-C100 Manufacturer EXBIO Category Antikörper Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen A KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP70 Product Description The ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase pr Clonality Monoclonal Concentration 1 mg/ml Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Article Name	Anti-ZAP70 Purified, Clone: [1E7.2], Monoclonal
Alternative Catalog Number EXB-11-665-C100 Manufacturer EXBIO Category Antikörper Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen A KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP70 The ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase pr Clonality Monoclonal Concentration 1 mg/ml Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Biozol Catalog Number	EXB-11-665-C100
ManufacturerEXBIOCategoryAntikörperApplicationFC, ICC, IP, WBSpecies ReactivityHuman, MouseImmunogenA KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP70Product DescriptionThe ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase prClonalityMonoclonalConcentration1 mg/mlClone Designation[1E7.2]IsotypeMouse IgG1 kappa	Supplier Catalog Number	11-665-C100
Category Antikörper Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen A KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP70 The ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase pr Clonality Monoclonal Concentration 1 mg/ml Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Alternative Catalog Number	EXB-11-665-C100
Application FC, ICC, IP, WB Species Reactivity Human, Mouse Immunogen A KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP70 The ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase pr Clonality Monoclonal Concentration 1 mg/ml Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Manufacturer	EXBIO
Species Reactivity Human, Mouse A KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP70 The ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase pr Clonality Monoclonal Concentration 1 mg/ml Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Category	Antikörper
Immunogen A KLH-coupled peptide corresponding to amino acids 282-307 of human ZAP70 The ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase pr Clonality Monoclonal Concentration 1 mg/ml Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Application	FC, ICC, IP, WB
Product Description The ZAP70 (zeta-associated protein of 70 kDa) tyrosine kinase was identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase pr Clonality Monoclonal Concentration 1 mg/ml Clone Designation [1E7.2] Mouse lgG1 kappa	Species Reactivity	Human, Mouse
Product Description identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR stimulation. ZAP70 is a Syk family tyrosine kinase pr Clonality Monoclonal Concentration 1 mg/ml Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Immunogen	
Concentration 1 mg/ml Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Product Description	identified as a tyrosine phosphoprotein that associates with TCR zeta subunit and undergoes tyrosine phosphorylation following TCR
Clone Designation [1E7.2] Isotype Mouse IgG1 kappa	Clonality	Monoclonal
Isotype Mouse IgG1 kappa	Concentration	1 mg/ml
	Clone Designation	[1E7.2]
Buffer Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	Isotype	Mouse IgG1 kappa
	Buffer	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage 2°C to 8°C	Storage	2°C to 8°C

Target	ZAP70
Antibody Type	Monoclonal Antibody
Application Notes	Flow cytometry: Recommended dilution for purified antibody: 1 $\mu g/ml$, intracellular staining. Recommended protocol for primary antibody conjugates using ADG Fix and Perm kit: 1) Add 50 μl peripheral blood to a 5 ml FACS tube, 2) Proceed surface staining w