

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD8 APC-Cy(TM)7, Clone: [MEM-31], APC/Cy7, Monoclonal EXB-T4-207-T100

| | |
|----------------------------|---|
| Article Name | Anti-Hu CD8 APC-Cy(TM)7, Clone: [MEM-31], APC/Cy7, Monoclonal |
| Biozol Catalog Number | EXB-T4-207-T100 |
| Supplier Catalog Number | T4-207-T100 |
| Alternative Catalog Number | EXB-T4-207-T100 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Human |
| Immunogen | Crude thymus membrane fraction. |
| Conjugation | APC/Cy7 |
| Product Description | The CD8 T cell coreceptor (monomer approx. 32-34 kDa) is expressed as alpha/beta heterodimer on majority of MHC I-restricted conventional T cells and thymocytes and as alpha/alpha homodimer on subsets of memory T cells, intraepithelial lymphocytes, N... |
| Clonality | Monoclonal |
| Clone Designation | [MEM-31] |
| Isotype | Mouse IgG2a |
| Buffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage | 2°C to 8°C |

| | |
|--------------------|--|
| Target | CD8 |
| Antibody Type | Monoclonal Antibody |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Application Notes | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |