

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD25 PE-Cy(TM)7, Clone: [MEM-181], PE/Cy7, Monoclonal EXB-T7-218-T100

| | |
|----------------------------|--|
| Article Name | Anti-Hu CD25 PE-Cy(TM)7, Clone: [MEM-181], PE/Cy7, Monoclonal |
| Biozol Catalog Number | EXB-T7-218-T100 |
| Supplier Catalog Number | T7-218-T100 |
| Alternative Catalog Number | EXB-T7-218-T100 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Human |
| Immunogen | PHA-activated peripheral blood leucocytes |
| Conjugation | PE/Cy7 |
| Product Description | CD25 (IL2Ralpha, Tac) is a ligand-binding alpha subunit of interleukin 2 receptor (IL2R). Together with beta and gamma subunit CD25 constitutes the high affinity IL2R, whereas CD25 alone serves as the low affinity IL2R. CD25 expression rapidly increas... |
| Clonality | Monoclonal |
| Clone Designation | [MEM-181] |
| Isotype | Mouse IgG1 |
| Buffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage | 2°C to 8°C |

| | |
|--------------------|--|
| Target | CD25 |
| Antibody Type | Monoclonal Antibody |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Application Notes | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |