

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Anti-Hu CD27 PE-Cy(TM)5, Clone: [LT27], PE/Cy5, Monoclonal EXB-T8-308-T100

|                            |   |
|----------------------------|---|
| Article Name               | Anti-Hu CD27 PE-Cy(TM)5, Clone: [LT27], PE/Cy5, Monoclonal  |
| Biozol Catalog Number      | EXB-T8-308-T100   |
| Supplier Catalog Number    | T8-308-T100   |
| Alternative Catalog Number | EXB-T8-308-T100   |
| Manufacturer               | EXBIO   |
| Category                   | Antikörper  |
| Application                | FC  |
| Species Reactivity         | Human   |
| Immunogen                  | Human peripheral blood lymphocytes  |
| Conjugation                | PE/Cy5  |
| Product Description        | CD27 is a transmembrane 55 kDa protein of the nerve growth factor-receptor family, expressed as a disulfide-linked homodimer on mature thymocytes, peripheral blood T cells and a subpopulation of B cells. Activation of T cells via TCR-CD3 complex resu... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [LT27]  |
| Isotype                    | Mouse IgG2a   |
| Buffer                     | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide   |
| Storage                    | 2°C to 8°C  |

|                    |  |
|--------------------|--|
| Target             | CD27   |
| Antibody Type      | Monoclonal Antibody  |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Application Notes  | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |