

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Hu CD16 PE-Cy(TM)5, Clone: [3G8], PE/Cy5, Monoclonal EXB-T8-646-T100

| | |
|----------------------------|---|
| Article Name | Anti-Hu CD16 PE-Cy(TM)5, Clone: [3G8], PE/Cy5, Monoclonal |
| Biozol Catalog Number | EXB-T8-646-T100 |
| Supplier Catalog Number | T8-646-T100 |
| Alternative Catalog Number | EXB-T8-646-T100 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Human, Primate |
| Immunogen | Human neutrophils |
| Conjugation | PE/Cy5 |
| Product Description | CD16 (FcgammaRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcgammaRIII is expressed in two forms - FcgammaRIII-A and -B. FcgammaRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T ... |
| Clonality | Monoclonal |
| Clone Designation | [3G8] |
| Isotype | Mouse IgG1 kappa |
| Buffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage | 2°C to 8°C |

| | |
|--------------------|--|
| Target | CD16 |
| Antibody Type | Monoclonal Antibody |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Application Notes | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |