

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-HLA-DR PE-Cy(TM)5, Clone: [L243], PE/Cy5, Monoclonal EXB-T8-690-T100

| | |
|----------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | Anti-HLA-DR PE-Cy(TM)5, Clone: [L243], PE/Cy5, Monoclonal |
| Biozol Catalog Number | EXB-T8-690-T100 |
| Supplier Catalog Number | T8-690-T100 |
| Alternative Catalog Number | EXB-T8-690-T100 |
| Manufacturer | EXBIO |
| Category | Antikörper |
| Application | FC |
| Species Reactivity | Canine, Human, Primate |
| Immunogen | Human B lymphocytes |
| Conjugation | PE/Cy5 |
| Product Description | HLA-DR, a member of MHC class II glycoproteins, that bind intracellularly processed peptides and present them to the Th cells, is composed of 36 kDa alpha chain and 27 kDa beta chain, both anchored in the plasma membrane. Together with other MHC II m... |
| Clonality | Monoclonal |
| Clone Designation | [L243] |
| Isotype | Mouse IgG2a kappa |
| Buffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage | 2°C to 8°C |

| | |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target | HLA-DR |
| Antibody Type | Monoclonal Antibody |
| Application Dilute | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Application Notes | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |