

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Monoclonal Antibody to Chromogranin A / CHGA (Neuroendocrine Marker)(Clone : CHGA/777), Clone: [CHGA/777], Mouse ABI-36-1055

Article Name	Monoclonal Antibody to Chromogranin A / CHGA (Neuroendocrine Marker)(Clone : CHGA/777), Clone: [CHGA/777], Mouse
Biozol Catalog Number	ABI-36-1055
Supplier Catalog Number	36-1055
Alternative Catalog Number	ABI-36-1055-100UG
Manufacturer	Abeomics
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human CHGA protein
Product Description	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para
Clonality	Monoclonal
Clone Designation	[CHGA/777]
Isotype	Mouse IgG1, kappa
NCBI	1113

UniProt	P10645
Buffer	$100~\mu g$ in $500~\mu l$ PBS containing $0.05\%BSA$ and 0.05% sodium azide. Sodium azide is highly toxic.
Purity	Affinity Chromatography
Form	Purified
Formula	$100~\mu g$ in $500~\mu l$ PBS containing $0.05\%BSA$ and 0.05% sodium azide. Sodium azide is highly toxic.
Target	Chromogranin A / CHGA (Neuroendocrine Marker)
Application Dilute	Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 min