

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Monoclonal Antibody to CD63 (Late Endosomes Marker)(Clone : MX-49.129.5), Clone: [MX-49.129.5], Mouse ABI-36-1960

| | |
|----------------------------|---|
| Article Name | Monoclonal Antibody to CD63 (Late Endosomes Marker)(Clone : MX-49.129.5), Clone: [MX-49.129.5], Mouse |
| Biozol Catalog Number | ABI-36-1960 |
| Supplier Catalog Number | 36-1960 |
| Alternative Catalog Number | ABI-36-1960-100UG |
| Manufacturer | Abeomics |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF, IHC, WB |
| Species Reactivity | Human, Mouse |
| Immunogen | Full length CD63 of human origin |
| Product Description | This MAb recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of MAb LAMP3/529. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and ... |
| Clonality | Monoclonal |
| Clone Designation | [MX-49.129.5] |
| Isotype | Mouse IgG1, kappa |
| NCBI | 967 |

| | |
|--------------------|--|
| UniProt | P08962 |
| Buffer | 100 µg in 500 µl PBS containing 0.05%BSA and 0.05% sodium azide. Sodium azide is highly toxic. |
| Purity | Affinity Chromatography |
| Form | Purified |
| Formula | 100 µg in 500 µl PBS containing 0.05%BSA and 0.05% sodium azide. Sodium azide is highly toxic. |
| Target | CD63 (Late Endosomes Marker) |
| Application Dilute | Flow Cytometry (0.5-1ug/million cells), Immunofluorescence (0.5-1ug/ml), Western Blot (0.5-1ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris wi |