

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Dystrophin Monoclonal Antibody (Clone: MANDYS8), Clone: [MANDYS8], Mouse ABI-39-1032

| | |
|----------------------------|---|
| Article Name | Anti-Dystrophin Monoclonal Antibody (Clone: MANDYS8), Clone: [MANDYS8], Mouse |
| Biozol Catalog Number | ABI-39-1032 |
| Supplier Catalog Number | 39-1032 |
| Alternative Catalog Number | ABI-39-1032-100UG,ABI-39-1032-100UG/VIAL |
| Manufacturer | Abeomics |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC-P, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant human dystrophin fragment. |
| Product Description | Dystrophin(DMD) gene has 79 exons spanning at least 2,300 kb(2.3 Mb). The C terminus of the dystrophin protein is encoded by a highly conserved, alternatively spliced region of the gene. beta-dystroglycan binding activity is expressed by the dystroph... |
| Clonality | Monoclonal |
| Clone Designation | [MANDYS8] |
| Isotype | Mouse IgG2b |
| NCBI | 24907 |

| | |
|--------------------|--|
| UniProt | P11530 |
| Buffer | Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN ₃ as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100 µg/ml. |
| Purity | Ascites |
| Formula | Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN ₃ as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100ug/ml. |
| Target | Dystrophin |
| Application Dilute | Western blot : 1-2µg/ml, Immunohistochemistry(Paraffin-embedded Section) : 2-4µg/ml |