

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Involucrin (Squamous Cell Terminal Differentiation Marker) Recombinant Rabbit Monoclonal Antibody (Clone:IVRN/2113R), Clone: [IVRN/2113R] ABI-12-1201-20

Article Name	Anti-Involucrin (Squamous Cell Terminal Differentiation Marker) Recombinant Rabbit Monoclonal Antibody (Clone:IVRN/2113R), Clone: [IVRN/2113R]
Biozol Catalog Number	ABI-12-1201-20
Supplier Catalog Number	12-1201-20
Alternative Catalog Number	ABI-12-1201-20-20UG
Manufacturer	Abeomics
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant full-length human IVL protein
Product Description	It recognizes a protein of 66kDa-170kDa, identified as involucrin. In Western blotting of cultured human keratinocytes, this MAb reacts with a 120kDa protein. It stains the involucrin in a variety of sizes: 170kDa in MCF-7 cells, a doublet of ~115kDa
Clonality	Monoclonal
Clone Designation	[IVRN/2113R]
Isotype	Rabbit IgG

NCBI	3713
UniProt	P07476
Buffer	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also available WITHOUT BSA & amp; azide at 1.0mg/ml.
Purity	Protein A/G
Form	Purified
Formula	$200~\mu g/ml$ of recombinant MAb purified by Protein A/G. Prepared in $10mM$ PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at $1.0mg/ml$.
Target	IVL
Application Dilute	Does not react with mouse.Flow Cytometry (0.5-1 μ g/million cells), ,Immunofluorescence (1-2 μ g/ml), ,Immunohistology (Formalin-fixed) (0.1-0.2 μ g/ml for 30 min at RT),(Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV