

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Anti-Thyroglobulin (Thyroidal Cell Marker) Recombinant Rabbit Monoclonal Antibody (Clone:TGB/1970R), Clone: [TGB/1970R] ABI-12-1251-20

Article Name	Anti-Thyroglobulin (Thyroidal Cell Marker) Recombinant Rabbit Monoclonal Antibody (Clone:TGB/1970R), Clone: [TGB/1970R]
Biozol Catalog Number	ABI-12-1251-20
Supplier Catalog Number	12-1251-20
Alternative Catalog Number	ABI-12-1251-20-20UG
Manufacturer	Abeomics
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Human thyroid follicular cells
Product Description	Thyroglobulin is a 660kDa dimeric pre-protein with mutiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo co
Clonality	Monoclonal
Clone Designation	[TGB/1970R]
Isotype	Rabbit IgG
NCBI	7038

UniProt	P01266
Buffer	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also available WITHOUT BSA & amp; azide at 1.0mg/ml.
Purity	Protein A/G
Form	Purified
Formula	$200~\mu g/ml$ of recombinant MAb purified by Protein A/G. Prepared in $10mM$ PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at $1.0mg/ml$.
Target	TG
Application Dilute	Immunohistochemistry (Formalin-fixed) (1-2µg/ml, 30 min at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes),