

Diagnostica Vertrieb GmbH, Leipziger Straße 4 85386 Eching, Germany Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99 E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat anti-Rabbit IgG (H&L), TRITC conjugated, Polyclonal AGR-AS10-1196

| Article Name | Goat anti-Rabbit IgG (H&L), TRITC conjugated, Polyclonal |
|----------------------------|---|
| Biozol Catalog Number | AGR-AS10-1196 |
| Supplier Catalog Number | AS10-1196 |
| Alternative Catalog Number | AGR-AS10-1196 |
| Manufacturer | Agrisera |
| Host | Goat |
| Category | Antikörper |
| Conjugation | tritc |
| Product Description | Goat anti-rabbit IgG (H&L), Rhodamine (TRITC-tetramethyl rhodamine-5-isothiocyanate) conjugated is a secondary antibody which binds to rabbit IgGs in immunological assays. TRITC has Amax = 550 nm, Emax = 570 nm. Antibodies are affinity purified using |
| Clonality | Polyclonal |
| Purity | Immunogen affinity purified IgG. |
| Form | Lyophilized |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | The optimal working dilution should be determined by the investigator, (ICC), immunohistochemistry (IHC) (Flow cyt) |
| Application Notes | For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily |