

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Donkey anti-Mouse IgG (H&L), FITC conjugated, Polyclonal AGR-AS10-1285

Article Name Donkey anti-Mouse IgG (H&L), FITC conjugated, Polyclonal Biozol Catalog Number AGR-AS10-1285 Supplier Catalog Number AS10-1285 Alternative Catalog Number AGR-AS10-1285 Manufacturer Agrisera Host Donkey Category Antikörper Conjugation fitc Donkey anti-mouse IgG (H&L), FITC (fluorescein-5-isothiocyanate) conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Supplier Catalog Number AS10-1285 Alternative Catalog Number AGR-AS10-1285 Manufacturer Agrisera Host Donkey Category Antikörper Conjugation fitc Donkey anti-mouse IgG (H&L), FITC (fluorescein-5-isothiocyanate) conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Alternative Catalog Number AGR-AS10-1285 Manufacturer Agrisera Host Donkey Category Antikörper Conjugation fitc Donkey anti-mouse IgG (H&L), FITC (fluorescein-5-isothiocyanate) conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Manufacturer Agrisera Donkey Category Antikörper Conjugation fitc Donkey anti-mouse IgG (H&L), FITC (fluorescein-5-isothiocyanate) conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Host Donkey Category Antikörper Conjugation fitc Donkey anti-mouse IgG (H&L), FITC (fluorescein-5-isothiocyanate) conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Category Antikörper Conjugation fitc Donkey anti-mouse IgG (H&L), FITC (fluorescein-5-isothiocyanate) conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Conjugation fitc Donkey anti-mouse IgG (H&L), FITC (fluorescein-5-isothiocyanate) conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Product Description Donkey anti-mouse IgG (H&L), FITC (fluorescein-5-isothiocyanate) conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Product Description conjugated is a secondary antibody which binds to mouse IgGs in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody
Clonality Polyclonal
Purity Immunogen affinity purified IgG.
Form Liquid
Antibody Type Polyclonal Antibody
Application Dilute 1 : 20-1 : 2000 (IHC), 1 : 50-1 : 5000 (ICC) (Flow cyt)