

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Donkey Antiserum anti-Rabbit IgG (Fc)-unconj., MinX none, Polyclonal DNA-SEC-182459

|                            |   |
|----------------------------|---|
| Article Name               | Donkey Antiserum anti-Rabbit IgG (Fc)-unconj., MinX none, Polyclonal  |
| Biozol Catalog Number      | DNA-SEC-182459  |
| Supplier Catalog Number    | SEC-182459  |
| Alternative Catalog Number | DNA-SEC-182459  |
| Manufacturer               | dianova   |
| Host                       | Donkey  |
| Category                   | Antikörper  |
| Application                | WB, IHC, ELISA  |
| Species Reactivity         | Rabbit  |
| Immunogen                  | Rabbit IgG F(c) fragment  |
| Conjugation                | Unconjugated  |
| Product Description        | Anti-Rabbit IgG F(c) generated in donkey is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of rabbit IgG and... |
| Clonality                  | Polyclonal  |
| Concentration              | 80 mg/mL  |
| Isotype                    | Ig  |
| Buffer                     | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |

|                    |  |
|--------------------|--|
| Purity             | This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-donkey serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observe |
| Formula            | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,Azide/BSA free   |
| Target             | Rabbit   |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | WB: 1:2,000 - 1:10,000   |
| Application Notes  | Antibody Anti-Rabbit IgG F(c) is suitable for immunoblotting (western or dot blot), ELISA, and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.  |