

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Mouse IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-182583

| Article Name | Goat IgG anti-Mouse IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated |
|----------------------------|--|
| Biozol Catalog Number | DNA-SEC-182583 |
| Supplier Catalog Number | SEC-182583 |
| Alternative Catalog Number | DNA-SEC-182583 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | WB, IHC, ELISA |
| Species Reactivity | Mouse |
| Immunogen | Anti-Mouse IgG was produced by repeated immunization with mouse IgG F(c) fragment in goat. |
| Conjugation | Unconjugated |
| Product Description | Anti-Mouse IgG F(c) antibody generated in goat detects specifically mouse IgG F(c). It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. R |
| Clonality | Polyclonal |
| Concentration | 10.0 mg/mL |
| Isotype | Ig |

| Buffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
|--------------------|--|
| Purity | Anti-Mouse IgG F(c) Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated |
| Formula | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Mouse |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Application Notes | Anti-Mouse IgG F(c) antibody has been assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution |