

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit F(ab)2 anti-Cat IgG (H+L)-HRPO, MinX none, Polyclonal DNA-SEC-182653

| Article Name               | Rabbit F(ab)2 anti-Cat IgG (H+L)-HRPO, MinX none, Polyclonal   |
|----------------------------|--|
| Biozol Catalog Number      | DNA-SEC-182653   |
| Supplier Catalog Number    | SEC-182653   |
| Alternative Catalog Number | DNA-SEC-182653   |
| Manufacturer               | dianova  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | WB, IHC, ELISA   |
| Species Reactivity         | Feline   |
| Immunogen                  | Cat IgG whole molecule   |
| Conjugation                | HRP  |
| Product Description        | F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper |
| Clonality                  | Polyclonal   |
| Concentration              | 10.0 mg/mL   |
| Isotype                    | Ig   |
| Buffer                     | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

| Purity             | This product is a F(ab)2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis a |
|--------------------|---|
| Formula            | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,Azide/BSA free  |
| Target             | Cat   |
| Antibody Type      | Polyclonal Antibody   |
| Application Dilute | WB: 1:1,000 - 1:10,000  |
| Application Notes  | This product has been assayed against 1.0 ug of Cat IgG in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:   |