

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 anti-Sheep IgG (H+L)-FITC, MinX none, Polyclonal DNA-SEC-182715

| Article Name | Rabbit F(ab)2 anti-Sheep IgG (H+L)-FITC, MinX none, Polyclonal |
|----------------------------|--|
| Biozol Catalog Number | DNA-SEC-182715 |
| Supplier Catalog Number | SEC-182715 |
| Alternative Catalog Number | DNA-SEC-182715 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IF, FLISA |
| Species Reactivity | Sheep |
| Immunogen | Sheep IgG whole molecule |
| Conjugation | FITC |
| Product Description | This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial pla |
| Clonality | Polyclonal |
| Concentration | 10.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| Purity | This product is a F(ab)2 fragment of an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by extensive dialysis a |
|--------------------|---|
| Formula | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal |
| Target | Sheep |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 2.8, IF Microscopy Dilution: 1:1,000 - 1:5,000 |
| Application Notes | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. |