

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Cat IgG (H+L)-Biotin, MinX none, Polyclonal DNA-SEC-182755

| | |
|----------------------------|---|
| Article Name | Goat IgG anti-Cat IgG (H+L)-Biotin, MinX none, Polyclonal |
| Biozol Catalog Number | DNA-SEC-182755 |
| Supplier Catalog Number | SEC-182755 |
| Alternative Catalog Number | DNA-SEC-182755 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | WB, IHC, ELISA |
| Species Reactivity | Feline |
| Immunogen | Cat IgG whole molecule |
| Conjugation | Biotin |
| Product Description | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglu... |
| Clonality | Polyclonal |
| Concentration | 2.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|--------------------|--|
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single p |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,Azide/BSA free |
| Target | Cat |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Application Notes | Secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods. |