

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit IgG anti-Chicken IgG (Fc)-HRPO, MinX none, Polyclonal DNA-SEC-182797

| Article Name               | Rabbit IgG anti-Chicken IgG (Fc)-HRPO, MinX none, Polyclonal  |
|----------------------------|---|
| Biozol Catalog Number      | DNA-SEC-182797  |
| Supplier Catalog Number    | SEC-182797  |
| Alternative Catalog Number | DNA-SEC-182797  |
| Manufacturer               | dianova   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | ELISA, WB   |
| Species Reactivity         | Gallus  |
| Immunogen                  | Chicken IgG F(c) fragment   |
| Conjugation                | HRP   |
| Product Description        | Anti-Chicken IgG F(c) Antibody detects specifically Chicken IgG F(c). It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind |
| Clonality                  | Polyclonal  |
| Concentration              | 1.5 mg/mL   |
| Isotype                    | Ig  |
| Buffer                     | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Chicken IgG, |
|--------------------|--|
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin   |
| Target             | Chicken  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | WB: 1:1,000 - 1:10,000   |
| Application Notes  | Anti-Chicken IgG F(c) Antibody has been tested in ELISA and Western Blot. Anti-Chicken IgG F(c) Antibody has been assayed against 1.0 ug of Chicken IgG in a standard capture ELISA using TMB as a substrate for 30 minutes at room temperature. A working dil |