

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Dog IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-182811

Article Name	Goat IgG anti-Dog IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated
Biozol Catalog Number	DNA-SEC-182811
Supplier Catalog Number	SEC-182811
Alternative Catalog Number	DNA-SEC-182811
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	DOT, ELISA
Species Reactivity	Canine
Immunogen	Dog IgG F(c) fragment
Conjugation	Unconjugated
Product Description	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglu
Clonality	Polyclonal
Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single p
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Dog
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Anti-Dog IgG F(c) antibody has been tested by ELISA and dot blot and is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.