

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Dog IgG (Fc)-Alk. Phos., MinX none, ALP, Polyclonal DNA-SEC-182817

Article Name	Goat IgG anti-Dog IgG (Fc)-Alk. Phos., MinX none, ALP, Polyclonal
Biozol Catalog Number	DNA-SEC-182817
Supplier Catalog Number	SEC-182817
Alternative Catalog Number	DNA-SEC-182817
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	WB, IHC, ELISA
Species Reactivity	Canine
Immunogen	Dog IgG F(c) fragment
Conjugation	ALP
Product Description	Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglu
Clonality	Polyclonal
Concentration	0.61 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0

Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single p
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target	Dog
Antibody Type	Polyclonal Antibody
Application Dilute	WB: 1:500 - 1:2,500
Application Notes	This product has been assayed against 1.0 ug of Dog IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:5,000 of the reconstitution concent